



Trabalho de Conclusão - Suplemento

Composition and structure of Heteroptera (Insecta: Hemiptera) assemblages along a vegetational gradient at Itacolomi State Park, southern Brazil

Thiago Coelho Nepomucena¹, Carlos Barreto^{1,2}  & Alexandre Silva de Paula¹ 

1. Universidade Federal de Ouro Preto, Ouro Preto, Minas Gerais, Brazil. 2. Algoma University, Sault Ste. Marie, Ontario, Canada.

Entomology Beginners, vol. 4: e053 (2023)

Table 2 - Supplement. Heteroptera recorded at Itacolomi State Park trails with UV light traps, 2010-2011. A1 represents the late succession community at Trilha do Forno I, A2 represents the early succession community at Trilha do Forno II, and A3 represents the rupestrian field at Trilha da Capela.

	Spring			Summer			Fall			Winter			N%
	A1	A2	A3	A1	A2	A3	A1	A2	A3	A1	A2	A3	
Lygaeidae													
<i>Cryphula</i> sp.		2											0.2%
<i>Neopamera</i> sp.	2	19	9		2	5							3.7%
<i>Nysius</i> sp.		1											0.1%
<i>Ozophora</i> sp.	1	4	5					6		1	3	1	2.1%
Pentatomidae													
<i>Banasa</i> sp.	10	2	1	8	1	4	40	17	26	15	2	7	13.3%
Halyini sp.	1	1			1	3				2			0.8%
Miridae													
<i>Allommatus</i> sp.			4			2			2			2	1.0%
<i>Carijoanus</i> sp.				1									0.1%
<i>Ceratocapsus</i> sp.	1		2	3		5	1	1	1				1.4%
<i>Creontiades</i> sp.	2	4	1	2		1			4	1		6	2.1%
<i>Cyclidolon</i> sp.		1	1										0.2%
<i>Derophthalma</i> sp.	3	2	5		1	2			3	2	1	5	2.4%
<i>Euchilocoris</i> sp.	2	2	7	6	10	13	23	20	118	3	4	13	22.1%
<i>Gauchocoris</i> sp.					3	6							0.9%
<i>Hyaliodocoris</i> sp.			2	1	1	1		1		3			0.9%
<i>Knightocoris</i> sp.	1							1			1	1	0.4%
<i>Lepidoxenus</i> sp.								1					0.1%
<i>Leprocapsus</i> sp.	1												0.1%
<i>Megalopsallus</i> sp.	2	6	11	13	8	12		2				1	5.5%
<i>Neostenotus</i> sp.	30	2	34	24	9	44	1	6	7	16	12	29	21.4%
<i>Paraxenus</i> sp.			1	2	2	6		1	8		1		2.1%
<i>Phytocoris</i> sp.	2	4	2	2	9	5			2	1	3	7	3.7%
<i>Phytocorisca</i> sp.	4	4	7	7	2	10	3		6	4		9	5.6%
<i>Poeas</i> sp.	1				1	3			1			1	0.7%
<i>Proba</i> sp.	6	9	3	5	4	3		7	4		1	1	4.3%

To be continued...

Table 2 - Supplement. Continue...

	Spring			Summer			Fall			Winter			N%
	A1	A2	A3	A1	A2	A3	A1	A2	A3	A1	A2	A3	
<i>Pseudocarnus</i> sp.				3			1						0.4%
<i>Taedia</i> sp.	1												0.1%
<i>Tropidosteptes</i> sp.				3		1				1	1		0.6%
Reduviidae													
Ectrichiinae sp.			1										0.1%
Emesinae sp.	1		2	4				1		1			0.9%
<i>Oncerotrachelus</i> sp.	1	1	1						1				0.4%
<i>Rasahus</i> sp.			11										1.1%
<i>Repipta</i> sp.	1		1										0.2%
<i>Sirthena</i> sp.	1												0.1%
<i>Stenopda</i> sp.	1			1									0.2%
<i>Zelurus</i> sp.			1										0.1%
Rhopalidae													
<i>Jadera</i> sp.	1	1	1										0.3%
Cydnidae													
<i>Pangeaus</i> sp.			1	1							1		0.3%
Total	76	65	114	86	54	126	69	64	183	50	30	83	100%